

# Localizing Interference during Naming: Convergent Neuroimaging and Neuropsychological Evidence for the Function of Broca's Area

Tatiana T. Schnur<sup>\*1</sup>, Myrna F. Schwartz<sup>2</sup>, Daniel Y. Kimberg<sup>3</sup>, Elizabeth Hirshorn<sup>4</sup>, H. Branch  
Coslett<sup>3</sup>, and Sharon L. Thompson-Schill<sup>3</sup>

<sup>1</sup> Rice University, Houston, TX USA

<sup>2</sup> Moss Rehabilitation Research Institute, Albert Einstein Healthcare Network, Philadelphia USA

<sup>3</sup> University of Pennsylvania, Philadelphia USA

<sup>4</sup> University of Rochester, Rochester NY USA

Classification: Biological Sciences, Neurosciences

Manuscript Information: 24 pages, 2 figures

In press, PNAS

\*Address for correspondence:

Tatiana T. Schnur

Rice University

Department of Psychology MS 25

6100 Main Street, Houston TX

e-mail: [ttschnur@rice.edu](mailto:ttschnur@rice.edu)

## Abstract

In order to produce a word, the intended word must be selected from a competing set of other words. In other domains where competition affects the selection process, the left inferior frontal gyrus (LIFG) responds to competition among incompatible representations. The aim of this study was to test whether the LIFG is necessary for resolution of competition in word production. Using a novel methodological approach applying the same rigorous analytic methods to neuropsychological data as is done with neuroimaging data, we compared brain activation patterns in normal speakers (using fMRI) with the results of lesion-deficit correlations in aphasic speakers who performed the same word production task designed to elicit competition during lexical selection. The degree of activation of the LIFG in normal speakers and damage to the LIFG in aphasic speakers was associated with performance on the production task. These convergent findings provide strong support for the hypothesis that the region of cortex commonly known as Broca's area (i.e., the posterior LIFG) serves to bias competitive interactions during language production.

In 1861, Paul Broca wrote “somewhere in these [frontal] lobes, one or several convolutions holds under their dependence one of the elements essential to the complex phenomenon of speech.” In the ensuing century and a half, investigations of the psychological and neural characterization of the “phenomenon of speech” have flourished. Here we unite the principal method of Broca’s day – the assessment of the relation between lesion location and cognitive impairments – with the primary human neuroscientific tool of the modern era – functional magnetic resonance imaging (fMRI) – in a rigorous evaluation of one putative element of speech: conflict resolution.

The need to resolve conflict during speech production is not self-evident; producing speech can feel spontaneous and easy. However every word produced is susceptible to error. The analysis of speech errors – both the relatively infrequent ones that are made by normal speakers and those that occur with much greater frequency in patients with acquired language disorders – has been a rich source of information about the speech production process. Such analyses have revealed that word selection during production is a naturally competitive process, determined by the relative degree of support for (i.e., activation of) a set of candidate words (1-3). The question addressed in this paper is whether a region of the frontal lobes commonly referred to as Broca’s area (i.e., the posterior portion of the left inferior frontal gyrus [LIFG]), an area implicated in controlled memory retrieval (4), multiple aspects of language processing (5-8), and competition among linguistic and non-linguistic representations (9, 10), is necessary for the resolution of conflict among competing lexical representations during word production. We posed this question of both normal and impaired speakers, using a word production task that isolates competition during lexical selection. In Experiment 1, we measured normal speakers’ brain

activations using functional neuroimaging (fMRI). In Experiment 2, we mapped performance deficits to lesion locations in participants with aphasia.

Our approach has three major methodological strengths. First, we used a well-studied speech production paradigm known to create competition during word selection (described in more detail below). Second, we explored both directions of the mapping between brain and behavior, namely, the effects of competitive speech production on the LIFG (with fMRI) and the effects of the LIFG on competitive speech production (with lesion-deficit analyses). Both “directions” of inference are necessary to understand a system with a many-to-many mapping of structure and function (cf. 11). Lastly, our lesion-deficit analysis combined a region-based approach with a statistical innovation in voxel-based lesion-symptom mapping (12) in order to search for relations with adequate specificity (low type I error) and sensitivity (low type II error).

#### Experiment 1 – fMRI analysis of interference effects on normal word production

Word selection during production is considered a naturally competitive and automatic process. The selection of a word is determined by the word with the highest activation level in comparison to other activated words (1-3). We manipulated lexical competition during simple picture naming by varying the context in which the pictures to be named appear, with successive trials depicting either semantically related items (semantically blocked context: e.g., truck, car, bike...) or mixed-category items (mixed context: e.g., truck, foot, dog...) named over four cycles. This blocking manipulation, referred to as the “blocked naming paradigm”, results in longer naming latencies to an item appearing in a semantic block than to the same item in a mixed block. This blocking effect has been attributed to competition among lexical items that are simultaneously activated due to their semantic relatedness (13-19), making it ideally suited for our purposes. This simple picture naming task offers advantages over other tasks in that the same

items are named in both high and low competition conditions, error rates are low, it is a pure production task (e.g., picture naming only), effects are robust across participants, and it is well studied in both unimpaired and aphasic populations (13-21).

Sixteen healthy volunteers named the pictures aloud in this blocked naming paradigm while undergoing BOLD imaging. As a control for effects of semantic blocking beyond increased competition (e.g., phonological and/or articulatory processes), we compared semantic blocked naming to a closely matched naming task, phonological blocked naming (naming pictures that sound the same vs. those mixed between phonological categories; cf. 14, 22). Blocking in each paradigm induces opposing behavioral effects (semantic interference, phonological facilitation; see Figure 1 A-B).

### Results and Discussion

We examined activity in both the LIFG and a left temporal region (consisting of the left superior and middle temporal gyri), the latter region's involvement suggested by magnetoencephalography (MEG) evidence in blocked naming (cf.17). The LIFG was significantly more active during semantically blocked compared to mixed naming  $t(1, 15) = 2.37, p = .03$ , but was unaffected by phonological blocking  $t < 1$ ; see Figure 1C. A direct comparison of blocking effect sizes for semantic and phonological experiments in the LIFG shows a significant greater involvement of LIFG in semantic blocked naming  $t(1,15) = 2.21, p = .04$ . The left temporal cortex was also significantly more active during semantically blocked compared to mixed naming  $t(1, 15) = 2.14, p = .04$ , an effect that did not obtain for phonologically blocked compared to mixed naming  $t(1,15) = 1.17, p > .20$ . However, a direct comparison between blocking effect size in the left temporal cortex for semantic and

phonological experiments shows no difference in activation between experiments  $t(1, 15) = 1.08, p = .29$ .

We also examined the right inferior frontal gyrus as an analogous region of interest (ROI) to the LIFG, and the anterior cingulate cortex because of its role in conflict monitoring in other tasks (24). Neither the anterior cingulate cortex nor the right inferior frontal gyrus was significantly more active in blocked compared to mixed naming in either the semantic or phonological paradigms.

To examine the specificity of the finding from the ROI analyses, we conducted a whole brain group analysis to see activation across the whole brain. The whole brain analysis supports the ROI analysis: the LIFG, specifically the pars triangularis in the posterior portion of the LIFG, and the left middle temporal gyrus were significantly more active in semantic blocked naming compared to phonological blocked naming. The whole brain analysis also revealed that the frontal involvement extended to the left middle frontal gyrus, and additional loci included the right superior temporal gyrus, the insula and lateral globus pallidus (see supporting information (SI) and Table S1).

Error rates across all conditions were low (semantically blocked: 1.2%; semantically mixed: 1.3%; phonologically blocked: 1.1%; phonologically mixed: 1.4%) but sufficiently variable to permit a correlational analysis of individual differences in psychological and neural blocking effects. Errors in semantic blocked naming are attributed to the misselection and production of semantically related competitors (18, 25). Thus, we assumed that a participant's error rate is an index of increased competition during naming. We correlated the number of errors in the semantic blocked condition with the activity in the LIFG, left temporal, right inferior frontal gyrus, and anterior cingulate ROIs. As depicted in Figure 1D, individuals with a

large LIFG response to semantic blocking tended to make more naming errors in the blocked condition  $r = .76, p < .0001$ ; this correlation was not found in the left temporal cortex  $r = .20, p = .46$  (Figure 1E) or the other regions,  $r$ 's  $< .30, p$ 's  $> .30$ . Further, the LIFG correlation was significantly different than the correlations in the other ROIs (Steigler's  $Z$ 's  $> 2.8, p$ 's  $< .01$ ). To rule out the possibility that the LIFG responds as an "error detector", we also correlated activity in the LIFG with the number of errors produced during the other three naming conditions (phonological blocked, semantic and phonological mixed). We found no significant correlations  $r$ 's  $< .15, p$ 's  $> .70$ . These analyses suggest that the LIFG responds to competition as measured by selection difficulty (e.g., erroneous naming) during blocked naming.

In summary, both the LIFG and left temporal cortex responded to the semantic blocking manipulation during speech production. This pattern was specific to semantic blocking (i.e., dissociated from phonological blocking) only in the LIFG. Additionally, the magnitude of the semantic blocking effect in the LIFG (but not left temporal cortex) correlated with the number of errors produced. We interpret these neural blocking effects as the result of increased demands to resolve conflict among competing lexical items made active by the blocking manipulation.

#### Experiment 2 – Lesion analysis of interference effects on impaired word production

In order to assess whether the LIFG is not only involved but also necessary to resolve competition during word production, we mapped lesions in a subset of patients reported in Schnur et al.'s (18) semantic blocked naming experiment (Experiment 2). In 18 patients with post-stroke aphasia, Schnur et al. observed a significant semantic blocking effect (phonological blocking was not manipulated), measured in errors; the magnitude of the blocking effect increased across naming cycles. These effects were significantly larger in patients diagnosed with Broca's aphasia, a syndrome characterized by low fluency and agrammatic speech and

associated with anterior damage, compared to patients with different aphasia syndromes.

However, damage to Broca's area (i.e., the posterior LIFG) is neither necessary nor sufficient to produce Broca's aphasia, so these results do not elucidate the putative role of this region in speech production. Thus, in Experiment 2 we conducted an anatomical study with 12 of 18 patients from Schnur et al. to definitively assess whether lesion location, specifically the LIFG, predicts impairment in resolving competition during naming. The only criterion for participation was the ability and willingness to undergo neuroanatomical scans (12 of 18 patients qualified); behavioral performance varied. Based on Schnur et al.'s findings, we predicted that damage to the LIFG would be associated with an increase in interference at later naming cycles (i.e., as competition among names increases but the capability to resolve the competition is impaired), while those with lesions outside the LIFG might show an overall interference effect, but one that did not increase at later cycles (i.e., competition among names increases but is resolved by an intact LIFG). Because these deficit analyses do not suffer from the temporal limitations of BOLD data, in this experiment we can specifically test the association between LIFG and the growth of interference over temporal repetitions.

### Results and Discussion

We performed a ROI analysis in which we assessed whether the percent damage in either the LIFG or the left temporal cortex predicted performance during naming in the blocked naming task (specifically, blocking growth -- the increase in the blocking effect across naming cycles). As the relation between degree of damage and behavior is not particularly well characterized by a linear function, we computed linear contrasts of the condition by cycle effect, to classify each patient as having either a large growth effect (i.e., only positive increases in blocking interference across cycles,  $f > 1$ ; mean = 2.61,  $n = 7$ ) or a small growth effect (e.g., no increase in



blocking interference across cycles,  $f < 1$ ; mean = .27,  $n = 5$ ). Patients showed a range in behavioral performance ( $F$  values 0-3.95, see Table S2). Individuals with a large growth effect had a greater extent of damage in the LIFG than did those with a small growth effect,  $t(10) = 3.59$ ;  $p < .01$  (see Figure 2A). There was no difference between these two groups in the extent of damage in the left temporal ROI,  $t(10) = 1.44$ ;  $p > .15$ . These results support our hypothesis that the LIFG plays a specific role in the regulation of competition that emerges in the semantic blocking paradigm.

Correlational analyses support this conclusion. There was no correlation between the magnitude of a patient's growth effect ( $f$ -value) and that individual's age, months post-stroke, overall lesion-size, or percent damage to left temporal cortex (all  $p$ 's  $> .25$ ). In contrast, the correlation between percent damage to LIFG and the growth effect was marginally significant,  $r = 0.56$ ,  $p < .06$  (see Figure 2B).

To confirm results from the ROI analysis, and assess whether other (non-predicted) anatomical areas contributed to the behavioral deficit, we performed two whole brain analyses. Using the two patient groups described above (high and low growth effects), we compared the lesion distribution between these groups by creating a lesion subtraction map (see Figure 2C). The results show that the most common area of lesion overlap for the group of patients with a large growth effect is centered in the LIFG. This result was confirmed by the statistical comparison of the behavioral growth effect (again, the  $f$ -value) at every voxel as a function of that voxel's status (i.e., damaged versus not damaged) in a given individual. We used permutation testing, a non-parametric approach to significance testing that has been applied to solve the multiple comparison problem in functional imaging (statistical comparisons made at every voxel number over 60,000) (26). The advantages of permutation testing are multifold,

especially in the case of voxel-based statistical mapping (12). The only voxels that were reliably related to the growth effect (at a permutation-derived threshold of  $p < .05$ ) were located in the dorsal portion of the LIFG (see Figure 2D).

In summary, three different lesion-deficit analysis strategies converged on the result that damage in the LIFG is associated with growth of interference over cycles of semantically blocked naming. That is, although all aphasic speakers will tend to have difficulty with word production, by definition, their ability to resolve competition that arises in the course of language processing appears to depend on the integrity of the LIFG.

### General Discussion

We opened with a quotation from Paul Broca, in which he proposes a relation between some part of the left frontal cortex and some aspect of speech production. The two experiments presented here provide a detailed specification of both aspects of this putative relation: Our findings indicate that although both the LIFG and the left temporal cortex respond to a manipulation that increases conflict among lexical representations competing for selection during word production, only the LIFG is necessary for resolution of heightened competition during blocked naming. These results provide a new interpretation of speech errors in some aphasic patients (i.e., those who have damage to the LIFG) and add to the evidence that the prefrontal cortex functions more generally to regulate cognitive processing by biasing competitive interactions among incompatible representations (27).

We are preceded in our attempt to use the semantic blocking paradigm to understand the role of frontal cortex in speech production by two case studies. Patient FAS, whose language testing and SPECT studies were consistent with dysfunction of left frontal regions, made more errors in semantic blocked compared to mixed naming (20). The effects occurred only on the

output side (i.e., in naming, but not a comprehension version of the task) and only when naming required lexical selection from semantics (e.g., picture naming, not reading) suggesting the impairment occurred during production at a lexical level. Another patient, BM, also made more errors naming pictures when blocked by semantic category than when pictures were mixed between categories (21). Although lesion localization information was not available, BM presented with right hemiparesis and her language profile was consistent with transcortical motor aphasia – a left anterior aphasia presentation. Their resulting interpretation – that speakers with aphasia consistent with anterior damage have increased difficulty in producing words in situations of high competition – receives rigorous support from our analyses of a larger group of aphasic speakers in the present study. In so doing, we have also illustrated the great potential of voxel-based lesion-symptom mapping, and, unlike previous applications of these methods, we have done so in a modestly sized group of patients and with a procedure that provides adequate statistical sensitivity and specificity.

The ability to subject neuropsychological data to the same rigorous analytic methods as is done with neuroimaging data, and then to examine direct parallels between the two, stands to substantially increase our potential for understanding the complex mappings of psychological functions onto neural structures. As demonstrated here, this approach successfully localized the mechanism that resolves heightened competition during word production to the neuroanatomical substrate known as Broca's Area. Damage to this mechanism may explain the hesitant multiword speech evinced by those described as Broca's aphasics. Thus this finding opens an exciting line of research into how multiword speech is produced seemingly effortlessly, and why it can be so difficult for those with damage to the prefrontal cortex.

## Methods

### Experiment 1

Subjects. We collected fMRI data from 16 volunteers (5 males, 11 females; age range 18-33).

All were right-handed, native English speakers, none reported neurological or neuropsychological illnesses or were on medication. Participants gave informed consent in accordance with policies of the Institutional Review Board (IRB) of the University of Pennsylvania, Philadelphia PA and were paid for their participation.

Materials. Each of the 12 semantically-blocked sets comprised six achromatic line drawings (e.g., coat, dress, glove, hat, skirt, and sock) from the same semantic category (e.g. clothing; other categories were animals, appliances, body parts, food, furniture, nature, plants, roles, shapes, toys, utensils). The same pictures composed 12 semantically-mixed sets of six pictures from different categories; for details see (18). For the phonological blocking control, 72 pictures were grouped into 12 phonologically-blocked sets of six items that shared the same initial phoneme (e.g., bear, belt, bird, boat, bone, and boot) and phonologically-mixed sets of items with different onsets (phonological and semantic stimuli shared 16 pictures). We refer to a set of six semantic, phonological, or mixed experimental pictures named four times as a block. Scrambled versions of every picture were created for a baseline task (detection of a horizontal or vertical line in the image).

Procedure. Each session contained 12 imaging runs, each comprising eight blocks of stimuli (two semantic, two mixed, four scrambled baseline), alternating between conditions (e.g., semantic, baseline, mixed, baseline, semantic, baseline etc.). Half of the subjects completed six runs of the semantic blocking paradigm (semantic blocks and their mixed controls) followed by six runs of the phonological blocking paradigm (phonological blocks and their mixed controls);

the other half of the subjects performed the phonological paradigm followed by the semantic paradigm.

In each paradigm, pictures within a set were presented an equal number of times. Each set of six pictures was repeated four consecutive times (cycles) in a different order, forming a block of 24 pictures. Thus, each participant saw six pictures in each of 12 semantic/phonological and 12 mixed sets, cycled 4 times for a total of 576 trials in each paradigm. In the baseline task, 12 scrambled pictures were presented in each of 48 blocks for a total of 576 trials. Each trial contained a fixation (500 ms), a blank screen (200 ms), stimulus presentation (650 ms), and a blank response period (1150 ms). Throughout the experiment, participants were instructed to name pictures as quickly as possible without sacrificing accuracy. A session lasted approximately 2 hours.

Image Acquisition. Participants were scanned at the University of Pennsylvania using a 3 Tesla Siemens Trio scanner with a standard 8-channel head coil. For each participant, T1-weighted anatomical images were obtained at the beginning of the session using a 3D MPRAGE pulse sequence (TR=1620 ms, TE=3 ms, TI 950 ms, voxel size= 0.9766 X 0.9766 X 1 mm, matrix size 192 X 256 X 160) before T2\* weighted functional images were acquired. We acquired 152 sets of 16 interleaved, axial gradient echo, echoplanar images with TR=2.5 TE=30, 64x64 pixels in a 19.2 cm field of view, voxel size = 3mm X 3mm X 5mm for each run. The image acquisition period was 1000 ms with a 1500 ms gap (TR=2500 ms), to minimize movement artifacts (28) and to allow the experimenter to hear overt responses. The gap was positioned (300 ms post stimulus onset) so that no participant produced a response during image acquisition. Online prospective motion correction was performed with a PACE sequence.

Image Processing and Analysis. Functional images were sinc interpolated in time to correct for the fMRI acquisition sequence and were spatially smoothed with an 8mm FWHM Gaussian kernel. We used the general linear model as implemented in VoxBo ([www.voxbo.org](http://www.voxbo.org)) to analyze data. Analysis included an empirically derived 1/f noise model, filters removing low temporal frequencies (below .0048 Hz), and regressors to model global signal variations and between-scan differences (29). Covariates of no interest were created for each of the four cycles of each condition of the picture naming task (eg., semantic blocked 1, 2, etc.), each lasting 15 sec. Each stimulus condition, modeled as a boxcar function, was convolved with a canonical hemodynamic response function.

## Experiment 2

Subjects. We identified 12 individuals with chronic aphasia secondary to left hemisphere stroke (from a group initially reported by (18)) based on their ability and willingness to undergo neuroimaging procedures. All participants were right-handed, native speakers of English with an average age of 53 years (range 35 – 68), average education of 14 years (range 10 – 19), and average time from stroke of 63 months (range 10 – 175). All patients gave informed consent in accordance with the IRB of Albert Einstein Medical Center (behavioral experiment) and the University of Pennsylvania (neuroimaging study) See Table S2 for individuals' age, months post-onset, and aphasia classification.

Materials and Procedure. The stimuli were identical to those described above for the semantic blocking paradigm (patients did not perform the phonological blocking paradigm). Unlike subjects in the fMRI study, patients completed the task at their own pace, with a 5-sec response deadline. We varied the response-stimulus interval within subjects (one or five seconds). The first complete response before the 5-second deadline was taken as the picture-naming response.

The target name, correctly pronounced, was scored as correct. Anything else was scored as an error (see (18) for a complete account of the behavioral methods).

The behavioral measure of interest was the growth of interference across naming cycles. For each subject, we calculated the error rate for each picture when in a semantic block and when in a mixed block, by cycle, and then we computed linear contrasts of blocking effect across cycles. For each participant, this resulted in a linear contrast  $f$ -value that described the growth of interference over cycles (independent of overall magnitude). Individual behavioral performance was based on 1,150 trials.

Lesion analyses. For six participants, T1-weighted magnetic resonance image volumes were collected on a 3 Tesla Siemens scanner. Each volume consisted of 160 contiguous axial slices covering the entire brain (matrix size 192 x 256; 1 mm voxels). For six participants who were unable or unwilling to undergo an MRI scan, we obtained CT data in 16-slice scanners. CT scan volumes consisted of at least 44 contiguous axial slices covering the entire brain (matrix size 512 x 512; 2.5 – 3 mm slice thickness; in 1 case, 32 slices of 9 mm slice thickness and in another case 33 slices of 5 mm slice thickness).

We identified lesion boundaries on a standard MNI template using MRIcro (30) after reslicing the MNI template to match the angle of acquisition for each participant's scan. We matched each slice of the template to a slice in the participant's scan and manually drew the lesion contour onto the corresponding template on axial slices. With MRI data, this produced a contiguous 3D lesion volume. As CT scans provide fewer slices than the template, we manually interpolated between CT slices to render the lesion on each slice of the template. All lesion reconstructions were performed by the same experimenter; however, for five patients, a second experimenter repeated the procedure to allow for a reliability assessment. The degree of

reliability (mean percent volume difference =  $23 \pm 11$ ; and mean percent discrepant voxels =  $6 \pm 5$ ) (discrepant is defined as more than two voxels from the other manually drawn lesion volume), was comparable to the same measures of inter-rater reliability reported elsewhere (31).

Each participant's lesion volume was then resliced at the angle of the MNI template and transferred to a higher-resolution version of the template (1x1x1mm; available as ch2.img in MRIcro) using MRIcro's tri-linear interpolation function. This was necessary in order to use the Automated Anatomical Labeling (AAL) map in MRIcro (aal.img). Using the AAL map, we defined two anatomical ROIs: the LIFG (inferior orbitalis, triangularis, and opercularis), and a temporal lobe area (superior and middle temporal gyri; cf. 17).

Across the 12 patients, the range of overall damage and in each ROI was as follows: Overall lesion volume 41 – 231 cc; LIFG: 0 – 77%; left Temporal Cortex: 0 – 44%. Table S2 includes for each patient overall lesion volume, the proportion of lesion damage in the LIFG, left temporal cortex and their corresponding anatomical subdivisions in MRICRO (LIFG: triangularis, opercularis, and orbitalis; left temporal cortex: superior and middle temporal gyri; based on (32)).

To further assess the specificity of the relation between LIFG damage and interference during word production, we performed two whole brain analyses. First, we compared the distribution of brain lesions in an overlay subtraction analysis, which revealed anatomical lesions unique to the high-growth interference group in comparison to the low-growth interference group (as seen elsewhere, e.g., 33-34). The advantage of this method is that lesions common to both groups are not revealed, only lesions different between groups. The disadvantage of this analysis is that participants must be divided into dichotomous groups, neglecting the reality that behavior is on a continuum (35). To address this concern, and more importantly to give a statistical



evaluation of the relationship between behavior and brain damage across the whole brain at every voxel (cf. 36), we used a nonparametric test, a permutation analysis (12, 26). See Figure 2D.

### Acknowledgements

This research was funded by grants from the National Institutes of Health: HD007425 (TTS); R01 DC000191 (MFS), MH60414 and MH067008 (STS). We gratefully acknowledge E. Lee's and A. Brecher's contribution to the data collection and analysis. We benefited from helpful discussions with Gary Dell. An analysis in this study was presented at the Academy of Aphasia, October 2005 in Amsterdam, Netherlands (Schnur TT, Lee E, Coslett HB, Schwartz MF, Thompson-Schill SL *Brain Lang* 95: 12 – 13).

## References

1. Dell GS (1986) A spreading-activation theory of retrieval in sentence production. *Psychol Rev* 93:283-321.
2. Levelt WJM, Roelofs A, Meyer AS (1999) A theory of lexical access in speech production. *Behav Brain Sciences* 22:1-75.
3. Roelofs A (1992) A spreading-activation theory of lemma retrieval in speaking. *Cognition* 42:107-142.
4. Badre D, Wagner AD (2007) Left ventrolateral prefrontal cortex and the cognitive control of memory. *Neuropsychologia* 45:2883–2901.
5. Grodzinsky Y, Santi A (in press) The battle for Broca's region. *Trends in Cog Sci*.
6. Hagoort P (2005) On Broca, brain, and binding: a new framework. *Trends in Cog Sci* 9:416-423.
7. Indefrey P, Levelt WJM (2004) The spatial and temporal signatures of word production components. *Cognition* 92:101-144.
8. Martin RC (2003) Language processing: Functional organization and neuroanatomical basis. *Annu. Rev. Psychol.* 54:55–89.
9. Thompson-Schill SL, D'Esposito M, Aguirre GK, Farah MJ (1997) Role of left inferior prefrontal cortex in retrieval of semantic knowledge: A reevaluation. *Proc Natl Acad Sci USA* 94:14792-14797.
10. Thompson-Schill SL et al. (1998) Verb generation in patients with focal frontal lesions: A neuropsychological test of neuroimaging findings. *Proc Natl Acad Sci USA* 95:15855-15860.

11. Sarter M, Berntson GG, Cacioppo JT (1996) Brain imaging and cognitive neuroscience: Toward strong inference in attributing function to structure. *American Psychologist* 51:13–21.
12. Kimberg DY, Coslett HB, Schwartz MF (2007) Power in voxel-based lesion-symptom mapping. *J Cogn Neurosci*, 19:1067-1080.
13. Belke E, Meyer AS, Damian MF (2005) Refractory effects in picture naming as assessed in a semantic blocking paradigm. *Quart J Expl Psychol Sec A Human Exp Psychol* 58:667-692.
14. Damian MF (2003) Locus of semantic interference in picture–word interference tasks. *Psych Bull Rev* 10:111-117.
15. Damian MF, Als LC (2005) Long-lasting semantic context effects in the spoken production of object names. *J Exp Psychol Learn Mem Cognit* 31:1372-1384.
16. Damian MF, Vigliocco G, Levelt WJM (2001) Effects of semantic context in the naming of pictures and words. *Cognition* 81:B77-B86.
17. Maess B, Friederici AD, Damian M, Meyer AS, Levelt WJM (2002) Semantic category interference in overt picture naming: Sharpening current density localization by PCA. *J Cogn Neurosci* 14:455-462.
18. Schnur TT, Schwartz MF, Brecher A, Hodgson C (2006) Semantic interference during blocked-cyclic naming: Evidence from aphasia. *J Mem Lang* 54:199-227.
19. Vigliocco G, Vinson DP, Damian MF, Levelt WJM (2002) Semantic distance effects on object and action naming. *Cognition* 85:B61-B69.
20. McCarthy RA, Kartsounis LD (2000) Wobbly words: Refractory anomia with preserved semantics. *Neurocase* 6:487-497.

21. Wilshire CE, McCarthy RA (2002) Evidence for a context-sensitive word retrieval disorder in a case of nonfluent aphasia. *Cognit Neuropsychol* 19:165-186.
22. Roelofs A (1999) Phonological segments and features as planning units in speech production. *Cognition* 14:173-200.
23. Buckner RL, et al. (1998) Functional-anatomic correlates of object priming in humans revealed by rapid presentation event-related fMRI. *Neuron* 20:285-296.
24. Botvinick MM, Braver TS, Barch DM, Carter CS, Cohen JD (2001) Conflict monitoring and cognitive control. *Psychol Rev* 108:624-652.
25. Vitkovitch M, Humphreys GW (1991) Perseverant responding in speeded naming of pictures- it's in the links. *J Exp Psychol Learn Mem Cogn* 17:664-680.
26. Nichols TE, Holmes AP (2002) Nonparametric permutation tests for functional neuroimaging: A primer with examples. *Hum Brain Mapp* 15:1-25.
27. Thompson-Schill SL, Bedny M, Goldberg RF (2005) The frontal lobes and the regulation of mental activity. *Curr Op Neurobiol* 15:219-224.
28. Huang J, Carr TH, Cao Y (2002) Comparing cortical activations for silent and overt speech using event-related fMRI. *Hum Brain Mapp* 15:39-53.
29. Aguirre GK, Zarahn E, D'Esposito M (1997) Empirical analyses of BOLD fMRI statistics.2. Spatially smoothed data collected under null-hypothesis and experimental conditions. *Neuroimage* 5: 199-212.
30. Rorden C, Brett M (2000) Stereotaxic display of brain lesions. *Behav Neurol* 12:191-200.
31. Fiez JA, Damasio H, Grabowski TJ (2000) Lesion segmentation and manual warping to a reference brain: Intra- and interobserver reliability. *Hum Brain Mapp* 9:192-211.

32. Tzourio-Mazoyer N et al. (2002) Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15:273-289.
33. Berti A et al. (2005) Shared cortical anatomy for motor, awareness and motor control. *Science* 309:488–491.
34. Karnath HO, Berger MF, Kuker W, Rorden C (2004) The anatomy of spatial neglect based on voxelwise statistical analysis: A study of 140 patients. *Cereb Cortex* 14:1164-1172.
35. Wu DH, Waller S, Chatterjee A (2007) The functional neuroanatomy of thematic role and locative relational knowledge. *J Cogn Neurosci* 19:1542-1555.
36. Bates E et al. (2003) Voxel based lesion-symptom mapping. *Nat Neurosci* 6: 448-450.

### Figure Legends

Figure 1. fMRI analysis of interference effects on normal word production. (A-B) Naming latencies in semantic and phonological blocked naming paradigms (error bars are one standard error of the mean). Because naming latencies were not obtained during fMRI scanning, a separate group of 18 volunteers (8 males, 10 females; age range 19-34) participated in a behavioral session so that naming latencies could be used to verify the predicted effects of semantic and phonological blocking with these materials and design. (A) We observed a semantic blocking interference effect [ $F1$  and  $F2$   $p$ 's  $< .0001$ ] that increased across cycles [interaction  $F1$   $p < .05$ ,  $F2$   $p < .0001$ ]. (B) In contrast, phonological blocking facilitated naming latencies [ $F1$  and  $F2$   $p$ 's  $< .0001$ ]. (C) fMRI blocking effects (blocked - mixed) in semantic and phonological paradigms, in LIFG and left temporal regions of interest (ROI) (y-axis reflects activation differences indexed by beta values; error bars are 95% confidence intervals). Anatomical regions were defined in each subject by sulcal boundaries. Within each of these anatomical regions, the ROI was further constrained to include only those voxels that were more activated during the first cycle (repetition) of both semantic/phonological and mixed blocks compared to baseline. We used the first cycle to create the functional picture naming ROIs because repetition has been shown to decrease activations in the LIFG and temporal cortex (23). Within each functional-anatomical ROI, we compared activation (indexed by beta values) associated with the two naming conditions. (D) The relationship between the numbers of errors produced in semantic blocked naming (x-axis) and the difference in signal between semantically blocked naming and the baseline task as indexed by beta values (y-axis). This relationship was significant in the LIFG ROI (y-axis;  $r = .76$ ,  $p < .0001$ ), but not in the left temporal ROI ( $r = .20$ ,  $p = .46$ ).

Figure 2. Lesion analysis of interference effects on impaired word production. (A) In the ROI analyses of lesion-deficit associations, patients who exhibited a large (HIGH) growth of interference across cycles of blocked naming ( $n = 7$ ) exhibited a greater extent of damage to the LIFG (expressed as a percentage of ROI volume) than did patients with a small (LOW) growth effect ( $n = 5$ ),  $t(10) = 3.59$ ;  $p < .01$ . (B) The correlation ( $r = 0.56$ ) between the magnitude of the growth effect (individual  $f$ -value describing the linear increase in semantic blocking across cycles) and the extent of damage to the LIFG. This relationship was not obtained in the left temporal ROI ( $r = .20$   $p = .52$ ). (C) Results of a voxel-based comparison of the lesion locations of patients with a large or small growth effect. The subtraction overlay analysis reveals the number of lesioned voxels in one group that overlap in a location not shared by the other group, across the whole brain. Voxels colored yellow were damaged in 7 of the 7 patients with a large growth effect (100%) and 0 of the 5 patients with a small growth effect (0%); voxels colored orange were damaged in at least 6 of the patients with a large growth effect and in no more than 1 of the patients with a small growth effect. (D) The effect of damage on naming behavior, evaluated at every voxel in which at least one patient was represented. In order to control for Type I error (avoid false positives) across the 62,990 comparisons, while maintaining adequate sensitivity (controlling Type II error, or false negatives), we determined the critical threshold via permutation test. Across the brain, we randomly re-paired the voxels' associated lesion values and behavioral scores 1,000 times. For each permutation, the maximum statistic observed anywhere in the brain was recorded. The 95th percentile maximum statistic is the threshold at which, when our experimental hypothesis is false, we expect to see even a single more extreme value only 5% of the time. Using this threshold can therefore be said to control the false positive rate at a map-wise corrected alpha of 0.05 (26). This procedure naturally accounts for the non-

independence of the 62,990 voxels in this sample (because for example, many voxels clustered together in one lesioned region will all have identical values of 'lesioned'). Permutation testing provides the same kind of control as Bonferroni correction (i.e., control of the family-wise error rate), while taking the non-independence of the observations into account. The figure displays the only region containing voxels that surpassed this threshold ( $t > 5.4$ ): the dorsal region of the LIFG (BA 44).



Figure 1





Figure 2



*Whole brain group analysis*

We conducted a whole brain analysis to see activation across the whole brain using a liberal uncorrected threshold in order to maximize sensitivity in this exploratory analysis. In order to control for activity associated with general aspects of naming in blocked contexts (e.g., processes incurred by repeatedly naming potentially salient blocked picture-sets) we directly compared activity associated with the semantic blocking effect with activity associated with the phonological (blocking) facilitation effect. This direct comparison (semblocked – semmixed – (phonblocked – phonmixed)), reveals those areas that are significantly more involved in the semantic blocking effect, independent of any general processes that occur when naming pictures in this type of paradigm.

To facilitate collection of overt responses, we restricted the data acquisition period to one second (allowing a 1.5 sec gap for overt responses). This constrained our data acquisition to 16 slices per individual, centered on the frontal lobe. Acquisition of superior and inferior areas varied between individuals depending on cortex size. Consequently, after normalizing the data to a common Montreal Neurological Institute space, our analyses were limited to z planes –14 to +68.

All functional data were normalized to the 1mm Montreal Neurological Institute template using a set of affine transformations with smoothly non-linear deformations, as implemented in SPM2. Functional images were resampled into 3 mm isotropic voxels and spatially smoothed with an 8 mm FWHM Gaussian filter. Individual *t*-maps were calculated for contrasts of interest and then smoothed to 12mm FWHM, to facilitate averaging of between-subject differences in functional organization in addition to differences in anatomical organization, before being entered in the analysis. Areas of activation from the analysis were identified at an uncorrected significance level of  $p < .01$  (critical *t*-value  $> 2.6025$ ) and are listed in Table S1.

## Table Legends

### Table S1.

Whole brain group analysis: MNI coordinates significantly more involved in the semantic blocking effect compared to the phonological blocking effect (significant at the  $p < .01$  level).

### Table S2.

Information on each individual included in Experiment 2. Abbreviations in table headings: OVL (overall lesion volume), Prop (proportion of voxels in the region that are damaged), LIFG (left inferior frontal gyrus), Oper (pars opercularis), Orbit (pars orbitalis), Triang (pars triangularis), STL (superior temporal lobe), MTL (medial temporal lobe).

Table S1.

Hemisphere	Region	X	Y	Z	<i>t</i>
<i>Left</i>					
	Inferior Frontal Gyrus (BA 45)	-48	21	24	3.35
	Insula (BA 13)	-33	15	-9	3.60
	Middle Frontal Gyrus (BA 9)	-51	9	39	3.76
	Middle Temporal Gyrus (BA 39)	-45	-66	30	3.32
<i>Right</i>					
	Lateral Globus Pallidus	21	-9	3	3.71
	Superior Temporal Gyrus (BA 22)	60	-51	15	4.05
	Superior Temporal Gyrus (BA 39)	48	-54	18	3.31

Table S2

Growth of												
Aphasia			months	OLV	Prop							blocking
Classificatio		(post	1mm	Prop	Prop	Prop	Prop	Temporal	Prop	Prop	effect	
sub	n	age	onset)	(cc)	LIFG	Oper	Orbit	Triang	Cortex	STL	MTL	(f values)
DBu	Anomic	50	52	96	0.15	0.35	0.04	0.14	0.35	0.58	0.24	0.07
ND	Anomic	60	29	44	0.35	0.94	0	0.34	0.01	0.03	0	0.23
RK	Anomic	50	36	104	0	0	0	0	0	0	0	0.68
AB	Anomic	57	118	98	0.56	0.98	0.47	0.46	0.08	0.26	0	2.14
SA	Anomic	35	10	52	0.5	0.9	0.16	0.57	0	0	0	2.97
BS	Broca	41	22	196	0.49	0.98	0	0.62	0.26	0.69	0.06	0
CC	Broca	56	28	127	0.75	0.56	0.62	0.91	0.43	0.57	0.37	1.45
LN	Broca	60	135	146	0.47	0.82	0.31	0.44	0.34	0.54	0.24	1.51
DB	Broca	56	175	231	0.51	0.82	0.01	0.73	0.44	0.28	0.52	2.59
AS	Broca	68	82	163	0.77	0.75	0.74	0.8	0.32	0.49	0.23	3.72
LCW	Broca	39	49	140	0.39	0.55	0.01	0.59	0.24	0.03	0.34	3.95
JBr	Conduction	61	16	41	0.01	0.03	0	0	0	0	0	0.39